



Bagworm And Its Control

Bagworms live in curious bag-like cocoons that are camouflaged with bits of foliage from the host plant. The common bagworm, *Thyridopteryx ephemeraeformis* (Haworth), is an interesting caterpillar. The most commonly observed form of this pest is the spindle-shaped silk bag camouflaged with bits of foliage, bark and other debris (Fig. 1). Completed bags range from 1-1/2 to 2-1/2 inches long. The larva within the bag is brown or tan, mottled with black, and the bee-like adult males have clear wings and fur covered bodies. The females remain larva-like and do not emerge from the bag. The larva may stick its head and front legs out of the top of the bag to feed and move. When disturbed, the larva immediately pulls its head into the bag and holds the opening closed. Mature larvae may stay on their host plant or drag their bags some distance before firmly attaching the bag for transformation into the adult stage.



The bagworm occurs in the eastern United States from New England to Nebraska and south through Texas. The larvae seem to prefer arborvitae and red cedar but many other conifers and deciduous trees are attacked. These include: pine, spruce, cypress, juniper, willow, black locust, sycamore, apple, maple, elm, poplar, oak, and birch.

Bagworm feeding on juniper



Types of Damage

Bagworm females cannot fly and local populations can build rapidly when established on preferred hosts, especially arborvitae, cedar and juniper. Crowded larvae may eat the buds on these conifers causing branch dieback and open, dead areas. Excessive defoliation of these conifers may cause entire plant death during the following season. Moderate defoliation is unsightly. This pest rarely builds up large populations in forested areas.

The mature larvae usually attach their bags to a branch by wrapping extra silk, which does not decay rapidly. This band of silk may girdle the branch as it grows, resulting in dead branches several years later. Be sure to cut off this silk band when removing bags from a plant.



Adult Male Bagworm

Life Cycle and Habits

Bagworms have a single generation per year and overwinter as eggs inside the female bag. There may be 300-1000 eggs in a bag. The eggs delay hatching until late-May or early-June. As they hatch, the small blackish larvae crawl out the bottom of the bag and spin down on a strand of silk. These larvae on a string are often picked up by the wind and ballooned to nearby plants. When a suitable host plant is found the young larva immediately begins to form a new bag over its body. This bag is only about 1/8 inch long and is soon covered with sawdust-like fecal pellets. As the larva feeds and grows it enlarges the bag and begins to incorporate bits and pieces of plant material. By mid-August, the larvae are mature and they often move to a sturdy branch or other structure where they attach the bag firmly with a strong band of silk. The larvae orient themselves with their heads down and pupate. The female pupa looks much like the larva while the male forms characteristic wing pads and other adult-like structures. Within four weeks the males emerge and actively fly about in search of females. Mature females call by releasing a sex attractant pheromone. After mating the female literally mummifies around the egg mass, this remains in place until the next spring. Eggs overwinter inside the bag and hatch in late spring when trees have all their leaves. Larvae spin a silken sack covering their body, which they take with them as they feed. In September, full-grown larvae attach the bag to a twig with silk, and then pupate. After several days, the males emerge and fly to females, which remain in their bags for mating. Females lay eggs, and then die.



Bagworm at 20% maturity

Control Measures

Bagworms are difficult to control because they are often unnoticed until mature. Mature larvae will often pupate early if they detect pesticides on the plant foliage. Though there are a few known parasites and predators, they are often not adequate in urban habitats.

Option 1-Cultural Control-Mechanical Hand Picking If the bags are few in number and easily reached they may be picked off the plant and squashed. This can be done easily in the late fall when deciduous foliage has been dropped or the bits of plant material on the bags turn brown and can be easily located on evergreens. Be sure to cut the attachment silk band so that the branch will not be girdled in the future.

Option 2-Chemical Control-Insecticide Sprays Stomach insecticides are very useful for control of bagworms. Remember that the plant foliage is to be thoroughly covered because the larvae are protected from contact by being in the silk bag. Again, early sprays against young larvae are more effective than later applications.

Option 3-Chemical Control-Timed Sprays Using Degree-Day Emergence Using a base temperature of 14.4 degrees C (57.9 degrees F) the median first emergence is 380 DD base 14.4 degrees C (716 DD degrees F) and the median last emergence is 572 DD base 14.4 degrees C (1062 DD degrees F). Allow for an extra week of ballooning after the last emergence mark has been reached before applying Bt or an insecticide. A less precise degree-day model uses a 50 degrees F base which calculates last emergence around 900 DD units.

Information obtained through the Ohio State Extension Factsheet HYG-2149-90



Insect and Disease Fact Sheet Compliments of New Century



www.newcenturytree.com

1-877-79TREES