



Insect and Disease Fact Sheet Compliments of New Century



www.newcenturytree.com

1-877-79TREES

Fletcher Scale



Identification.

In the past, Lecanium (*Lecanium fletcheri*) spp. of scales found on *Taxus* have been assumed to be *L. fletcheri*. However, *Lecanium fletcheri* is reported to have only one generation per year; while a common *Lecanium* scale found on *Taxus* plants has two generations per year. In some places in Michigan and nearby a *Lecanium* spp. on *Taxus* has one generation per year.

Host Plants.

All *Taxus* species. Also, a similar scale is found on arborvitae.

Life Cycle.

The first generation crawlers emerge from June 10-30, settle on the needles and twigs and feed until August when they are nearly mature. A second generation of crawlers emerges in September. These young scales will mature in May.

Scouting.

Look for small (3-5 mm), brown hemispherical scales on inner foliage and twigs of *Taxus* plants. Young immature scales are even smaller, flat and clear.

Management.

Only apply insecticides to *Taxus* if the scale population is too high (greater than 15% of samples are infested). Parasites and predators regulate scale populations. Parasites are most effective along field borders. Biological control has good potential in production nurseries as well.

Apply.

Dursban or Talstar in early October if scales can be found on more than one of 10 leaf samples examined in August. If an unacceptable level of live scale is still found in April or early May, another application can be made at that time. Early July is the next best time to apply an insecticide. However, because of a second generation, a July spray without an October spray is risky.

Information obtained through Michigan State Extension, Landscape CAT Alert 61094006



Insect and Disease Fact Sheet Compliments of New Century



www.newcenturytree.com

1-877-79TREES